

Yunnan Baiyao Capsules Instruction Manual

Please read the instruction carefully and use it under the guidance of physician
It contains Aconiti Kusnezoffii Radix Cocta; it is contraindicated in pregnant women and individuals with an allergic reaction to Yunnan Baiyao; be sure to clean the wound surface before external use.

[Pharmaceutical Name] Yunnan Baiyao Capsules

[Ingredients] National Secret Formula. It contains Aconiti Kusnezoffii Radix Cocta, other ingredients are omitted.

[Description] This product is hard capsule and the content is powder from grayish yellow to light brownish yellow; it has a special aroma, slightly cool taste and numbness in the tongue. The insurance pills are red spherical or spheroid-like waterpill with a brown or tan profile, a slight odor and bitter taste.

[Function and Indication] Eliminating blood stasis and hemostatic, promoting blood flow and analgesic, detoxifying and subsiding swelling. It is used for treatment of traumatic injury, stagnated blood swelling and pain, spitting blood, hemoptysis, hematochezia, hemorrhoidal bleeding, metrorrhagia and metrostaxia, suppurative and pyogenic infections and soft tissue bruise, closed fracture, brochiectasis and hemoptysis of pulmonary tuberculosis, ulcerative bleeding and infective diseases on skin.

[Specification] Each capsule content 0.25g.

[Administration and Dosage] As to wounds by knife, bullet or traumatic injury, no matter how severe they have been, the capsules are administered with warm boiled water by the bleeding wounded; The wounded suffer from stagnated blood, swelling and pain and no bleeding wounded take the capsules with wine; For patients suffer from diseases of gynecopathy take capsules with wine, but for those suffer from menorrhagia, metrorrhagia take capsules with warm boiled water. For initial stage of venous sore, patient administer a dosage of 0.25g and take another portion of medicine powder to mix homogeneously with wine which is used to apply on infectious part; for suppurative sore, oral administration is needed only. For slight injury and other internal hemorrhage symptom, capsules can be administered orally.

Oral Administration: 1-2 capsules each time, 4 times daily (for children: 2 to 5 years old take 1/4 of dosage; 6 to 12 years old take 1/2 of dosage). For severe traumatic injury, a pill of Insurance pill can be administered at first, as to slight injury no Insurance pill will be needed.

[Adverse reaction] Monitoring data after marketing showed that the product has the following adverse reactions:

1. Gastro-intestinal system disorders: nausea, vomiting, abdominalgia, diarrhea, stomach discomfort, etc.
2. Respiratory system disorders: dyspnea, tachypnea, throat discomfort, etc.
3. Skin and appendages disorders: eyelid edema, skin redness, general intense itching, urticaria in the trunk and limbs and others, etc.
4. In addition, there are high fever, chills, dizziness, headache, palpitations, chest tightness, pseudomembranous colitis, acute gastritis, menstrual disorders, increased menstruation, anaphylactic shock, etc.

[Contraindication] 1. It is contraindicated in pregnant women.

2. It is contraindicated in individuals with an allergic reaction to Yunnan Baiyao.

3. It is contraindicated in defective or purulent skin and mucous membrane.

[Precautions]

1. Avoid fava beans, fish and cold sour food during medication and within one day after drug withdrawal.
2. Be sure to clean the skin before external use and only when malignant ulcer begins.
3. Be careful with allergic constitution and a history of drug allergy.
4. Use strictly according to the usage and dosage of the drug instructions, not suitable for long time and large area. It must be applied under the monitoring of doctors if necessary to use large doses of this medicine in clinic.
5. To discontinue immediately if adverse reactions or suspected adverse reactions occur (thoroughly cleaning the site of medication if external use), and go to the hospital for treatment.
6. Athletes should use carefully.
7. Note: The Insurance pill is placed in the middle of blisterpack.
8. It contains Aconiti Kusnezoffii Radix Cocta, please carefully read instruction manual and purchase and use it under the guidance of the physician.
9. To consult physician and pharmacist before using it if other drugs are being used.
10. Keep out reach of children.
11. When taking this medicine, it is not suitable to take Pinelliae Rhizoma, Trichosanthis Fructus, Trichosanthis Semen, Trichosanthis Pericaepium, Trichosanthis Radix, Fritillariae Cirrhosae Bulbus, Fritillariae Thunbergii Bulbus, Fritillariae ussuriensis Bulbus, Fritillariae Pallidiflorae Bulbus, Fritillariae Hupehensis Bulbus, Ampelopsis Radix, Bletillae Rhizoma and its preparations at the same time.

[Pharmacological & toxicological effects]

1. Hemostatic effects: Promoting the aggregation of platelet in rat and rabbit significantly, enhancing the activity percentage and expression of surface glycoprotein of platelets, shortening the coagulation time, bleeding time of wounds and prothrombin time in rats and rabbits, demonstrating significantly constriction of artery vessel strip of rabbit.
2. Effects of promoting blood flow to eliminate blood stasis: Inhibiting thrombogenesis in vein of rats, remitting disturbance of microcirculation induced by high molecular dextrose in rats and improving the situation of blood flow, accelerating blood flow in auricle microcirculation of rats.
3. Anti-inflammatory action: Demonstrating significant antagonizing effect to the inflammatory models in animals induced by inflammation genesis factors, such as, adjuvant, agar, foreign protein, inflammatory genesis chemicals and granuloma formation around cotton pellet.
4. Healing wounds effects: Promoting the production of (bFGF) (blast fibrocyte growth factor) and vessel endothelium growth factor (VEGF) in rats significantly, promoting the expression of (bFGF) in the field of operation in rats and granulation tissue proliferation. bFGF and VEGF may promote the production of vascular endothelial cell and blast fibrocyte thus accelerating the growth of blood vessels and connective tissue proliferation to achieve the effect of healing wounds.

[Storage] Keep airtight in a dry place.

[Package] Sealed in Aluminium-plastic-Aluminium bolt, 16 capsules per plate, 1 insurance pill.

[Validity] 5 years.

[Execution standard] The first edition of Chinese pharmacopoeia in 2020

[Approval document number] GYZ No. Z53020799.

[Marketing Authorisation Holders and Producer] Yunnan Baiyao Group Co., Ltd.

Address: Yunnan Baiyao Group Co., Ltd., Chenggong New District, Kunming, Yunnan Province, China.

Postal code: 650500

Phone number: 4001000538

Fax number: 0871-66226685

Registered address: Number 3686 in Baiyao Street, Chenggong District, Kunming City, Yunnan Province

Web site: www.yunnanbaiyao.com.cn



YUNNAN BAIYAO GROUP CO., LTD.